

A dichotomous key to the genus *Drosera* L. (Droseraceae)

by

Dr. Jan Schlauer, Zwischenstr. 11, D-60594 Frankfurt/Main, Germany
email: jan@pbc-ths1.pci.chemie.uni-tuebingen.de

Introduction

Since the world-wide monograph of the genus *Drosera* by Diels, (1906), many new species were described, and several treatments for local floras appeared. In order to be able to identify even wrongly labelled (or unlabelled) specimens in herbaria as well as in horticultural collections, and to provide field workers with a guidance, a dichotomous key was compiled from the data available by herbarium and literature study. As far as possible, all taxa described validly (and considered distinguishable) until present were included. The key is thought to reflect phylogenetic development to a certain degree, inasmuch as the taxa are probably of monophyletic origin (with the exception of *sects. Oosperma* and *Drosera*, which may be more closely related to each other than is evident in the key). The key does rely on morphological characters even if in some cases the taxa share additional (phytochemical or cytological) features, which cannot be examined in dried herbarium specimens.

The infrageneric rearrangement by Seine & Barthlott (1994) provides little new insight, and it suffers considerably from the omission of recent information, e.g. Kondo & Lavarack's important cytological work (1984) is ignored. The key by Marchant (1982) is misleading in important distinguishing features (e.g. stipules are not "usually absent or inconspicuous" in *subgen. Drosera*, not even in his version of this subgenus, and not even in the Australian representatives!). Substantial work has been performed on palynology (Takahashi & Sohma, 1982), and phytochemistry (Zenk, Fürbringer & Steglich, 1969; Culham & Gornall, 1994). A reference list of synonyms (at the rank of species or below) for the genus *Drosera* was already published in an earlier paper (Schlauer, 1987). A permanently updated version thereof is accessible via internet (http://www.hpl.hp.com/bot/cp_home; the Carnivorous Plant Homepage, maintained by Rick Walker).

The realignments and the key to this second largest genus of carnivorous plants (about 135 species recognized here, compared with 215 in *Utricularia*, Lentibulariaceae) are not meant to represent a final conclusion. These should rather be understood as a help and a starting point for the many enthusiasts as well as the quite numerous scientists interested in this fascinating group. Even if some of the new combinations proposed here are rather preliminary, it is preferred to give all taxa used in the key valid names (however, due to lack of suitable type material, this is impossible in one case discussed below). I am well aware of several remaining shortcomings and numerous unresolved problems but I hope this modest contribution may incite efforts to overcome these.

Infrageneric realignment of the genus *Drosera* L.

The infrageneric subdivision of the genus as presented in the key necessitates formal validation of some taxa. To this purpose, a nomenclatural conspectus of the taxa of *Drosera* above the rank of species, including the most important synonyms, is presented below. In this, some abbreviations are used:

T	Type
LT	Lectotype
S	Synonym(s)
BN	Basionym

Drosera L., Spec.Pl.ed.1:281 (1753)

T: **D.rotundifolia** L.

S: *Rossolis* (Tournef. ex) Adans., Fam.2:245 (1763) *nom.superfl.*

T: *R.rotundifolia* (L.) Adans. *nom.illeg.* = **D.rotundifolia** L.

Rorella (Hall.ex) Allioni, Fl.Pedem.2:88 (1785) *nom.superfl.*

T: *R.rotundifolia* (L.) Allioni *nom.illeg.* = **D.rotundifolia** L.

Esera Neck., Elem.Bot.2:160 (1791)

T: *E.cistiflora* (L.) Neck. = **D.cistiflora** L.

Adenopa Raf., Fl.Tellur.3:37 (1836)

T: *A.anglica* (Huds.) Raf. = **D.anglica** Huds.

Dismophyla Raf., l.c.:36

T: *Dismophyla binata* (Labill.) Raf. = **Drosera binata** Labill.

Filicirna Raf., l.c.:37

T: *F.filiformis* (Raf.) Raf. = **D.filiformis** Raf.

Sondera Lehm., Pugill.8:44 (1844)

T: *S.macrantha* Lehm. = **D.heterophylla** Lindl.

D.subgen.Thelocalyx (Planch.) Drude in Engl. & Prantl, Nat.Pflanzenfam.3:271 (1891)

BN: *D.sect.Thelocalyx* Planch., Ann.sci.nat.3.ser.9:92 (1848)

T: **D.burmannii** Vahl

S: *D.sect.Rorella* DC., Prodr.1:317 (1824) *p.p.*

The pentamerous gynoecium, known only in one other subgenus (viz. **Bryastrum**), is a sufficient reason to keep this distinct from the rest of the genus. The two species belonging here, one from tropical Asia and N Australia, the other from S America share so many (assumedly primitive) features that a phylogenetic position close to the origins of the genus may be supposed.

D.subgen.Arcturia (Planch.) Schlauer *stat. nov.*

BN: *D.sect.Arcturia* Planch., Ann.sci.nat.3.ser.9:91 (1848)

T: **D.arcturi** Hook.

S: *D.sect.Rorella* DC., Prodr.1:317 (1824) *p.p.*

D.sect.Drosera auct. non L.: Seine & Barthlott, Taxon 43:584 (1994) *p.p.*

D.sect.Psychophila auct. non Planch.: Diels, Pflanzenr.26:62 (1906) *p.p.*

The separation of this subgenus, native from SE Australia to New Zealand, from the rest of the genus is claimed here on the basis of exstipulate, sheathing leaf bases. As pollen seems to be different between the two species (Culham, Am.J.Bot.80 Suppl.6:142, 1993), the leaf characteristics may be of convergent nature to a certain degree, however. Contrary to Diels (1906), inclusion of **D.uniflora** here is not supported (cf. **D.subgen.D.sect.Ptycnostigma**).

D.subgen.Stelogyne (Diels) Schlauer *stat.nov.*

BN: *D.sect.Stelogyne* Diels, Pflanzenr.26:103 (1906)

T: **D.hamiltonii** C.R.P.Andrews

S: *D.sect.Drosera* auct. non L.: Seine & Barthlott, l.c. *p.p.*

The fusion of the styles in this monotypic Australian subgenus is such a unique feature that segregation at more than sectional level seems inevitable.

D.subgen.Meristocaulis (Maguire & Wurdack) Schlauer *stat.nov.*

BN: *D.sect.Meristocaulis* Maguire & Wurdack, Mem.NY Bot.Gard.9:332 (1957)

T: **D.meristocaulis** Maguire & Wurdack

A single species with numerous distinguishing features, the most important of which

being undivided styles. Apparently a rather ancient relict on the Neblina peak.

D.subgen.Regiae Seine & Barthlott, l.c.:586

T: **D.regia** Stephens

S: *D.sect.Psychophila* auct.non Planch: Stephens, Trans.Roy.Soc.S.Af.13:309
(1926) p.p.

D.sect.secundistyla Culham, Novon (in press)

D.ser.Eurossolis Diels in Engler & Prantl, Nat.Pflanzenfam.2.ed.17b:781

(1936) p.p.

Sufficient palynological reasons for subgeneric separation of this primitive S African species have been presented by Takahashi & Sohma (1982), already.

D.subgen.Coelophylla (Planch.) Schlauer stat. nov.

BN: *D.sect.Coelophylla* Planch., Ann.sci.nat.3.ser.9:93 (1848)

T: **D.glanduligera** Lehm.

The obviously primitive pollen type does not allow inclusion of this Australian species in **subgen.Drosera**.

D.subgen.Lasiocephala (Planch.) Schlauer stat. nov.

BN: *D.sect.Lasiocephala* Planch., Ann.sci.nat.3.ser.9:93 (1848)

T: **D.petiolaris** R.Br.

S: *D.sect.Rorella* DC., l.c.:317 p.p.

D.sect.Ergaleium DC., l.c.:319 p.p.

D.sect.Rossolis auct. non Planch.: Diels, (1906):62 p.p.

D.sect.Polypeltes Diels, (1906):62 p.p.

The most outstanding feature of this subgenus (native to tropical N Australia and New Guinea) is the subpeltate to peltate lamina, not known in any species of **subgen.Drosera** proper. The completely peltate lamina alone was sufficient for both Diels (1906) and Marchant (1982) to shift **D.banksii** (which was included here by Planchon, 1848) to **subgen.Ergaleium**. Kondo & Lavarack (1982) have shown by cytological similarity that this species is closest to the **D.petiolaris** complex. Morphological features (presence of stipules and absence of tuber) have led Seine & Barthlott (1994) to the same conclusion. Another important argument is style morphology.

Of all subgenera segregated here, this is the closest to **subgen.Drosera**, but in the present situation a separation seems favourable.

As the lamina margin of **D.neocaledonica** (endemic to New Caledonia) is continuous with the petiole margin, this species (formerly included here because of dubious stipule and indumentum features by Diels, 1906) should be shifted to **subgen.Drosera** (**sect.Oosperma**).

D.subgen.Drosera

S: *D.sect.Rorella* DC., l.c. p.p. nom. superfl. (cf. **D.subgen.Bryastrum**)

S: *D.subgen.Rorella* (DC.) Diels, (1906):92 nom. superfl.

(cf. **D.subgen.Bryastrum**)

D.subgen.D.sect.Prolifera C.White, Vict.Nat.57:94 (1940)

T: **D.prolifera** C.White

S: *D.sect.Drosera* auct. non L.: Seine & Barthlott, l.c.:584 p.p.

D.sect.Arachnopus auct. non Planch.: Diels, (1906):77 p.p.

This section includes not only **D.prolifera** but also **D.schizandra** and **D.adelae**. These three tropical N Australian species are considered more closely related to each other than is any of them to **D.indica**, which should be excluded from this section, therefore.

D.subgen.D.sect.Arachnopus Planch., l.c.:92

T: **D.indica** L.

S: *D.subgen.Arachnopus* (Planch.) Drude, l.c.:272

D.sect.Rorella DC., l.c. :319 p.p.

D.sect.Drosera auct.non L.: Seine & Barthlott, l.c. p.p.

A single paleotropic species.

D.subgen.D.sect.Ptycnostigma Planch.

BN: *D.sect.Ptycnostigma* Planch., Ann.sci.nat.3.ser.9:92 (1848)

LT: ***D.pauciflora*** Banks ex DC. (Seine & Barthlott, l.c.:585)

S: *D.sect.Rorella* DC., l.c. :317 p.p.

D.sect.Rossolis Planch., l.c.:92 p.p.

D.sect.Drosera auct. non L.: Seine & Barthlott, l.c. p.p.

D.sect.Psychophila Planch., l.c.:91

T: ***D.uniflora*** Willd.

D.subgen.Ptycnostigma (Planch.) Diels, (1906):62

BN: *D.sect.Ptycnostigma* Planch.

With the distinguishing features as considered significant here (reduced stipules, but rudiments often visible, frequently thickened roots as storage organs, a tendency to form rather large corollas with wide corolla lobes), the circumscription of this section is widened considerably. It now includes all species of ***subgen.Drosera*** with the stipules adnate to the petiole. A noteworthy rearrangement is the inclusion of the two American species ***D.brevifolia*** (formerly included in *sect.Drosera*) and especially ***D.uniflora*** (formerly grouped with or near what is considered another subgenus in this treatment, viz. *Arcturia*). The last mentioned species shows close morphological (incl. palynological) affinities with ***subgen.Drosera***, however, and it is felt that its placement here does more accurately reflect phylogenetic relationship.

D.subgen.D.sect.Oosperma Schlauer *sect.nov.*

Folia stipulis conspicuis obsita. Styli 3, basi bifurcati, deinde integri, stigma integrum vel nonnunquam bilobum, rarissime iterum divisum. Semina ellipsoidea ad ovoidea.

T: ***D.intermedia*** Hayne

S: *D.sect.Rorella* DC., l.c. :318 p.p.

D.sect.Rossolis Planch., l.c.:92 p.p.

D.ser.Eurossolis Diels, (1906):81 p.p.

This section is a possibly fairly inhomogeneous grouping of all species formerly included in *sect.Drosera*, but differing from ***D.rotundifolia*** and its allies by the seeds being ovoid rather than fusiform. Further research is necessary to elucidate the natural affinities of this section. In its present circumscription, this section is nearly as widespread as *sect.Drosera*.

D.subgen.D.sect.Drosera

S: *D.sect.Rorella* DC., l.c. :317 nom. superfl. (cf. ***D.subgen.Bryastrum***)

D.sect.Rossolis Planch., l.c. nom.superfl.

T: ***D.rotundifolia*** L.

D.ser.Eurossolis Diels (1906):81 nom.superfl.

T: ***D.rotundifolia*** L.

D.sect.Cripterisma Planch., l.c.

T: ***D.hilaris*** Cham. & Schlechtd.

D.sect.Vagae Drude, l.c.:271

T: ***D.capensis*** L.

A rather homogeneous grouping, and the only one which has reached a wide distribution on all continents with the exception of Antarctica (however with only few species in the northern hemisphere). Considerable range extensions at least of the

(nearly) circumboreal species must have occurred in rather recent evolutionary times. Some of the youngest species of the genus have to be sought here. The group itself is not necessarily an advanced one, however.

D.subgen.Bryastrum (Planch.). Schlauer *stat. nov.*

BN: *D.sect.Bryastrum* Planch., Ann.sci.nat.3.ser.9:94 (1848)

T: ***D.pygmaea*** DC.

S: *D.sect.Rorella* DC., l.c. p.p.

D.subgen.Rorella (DC.) Diels, (1906):81 p.p.

D.subgen.Rorella auct. non (DC.) Diels: N.Merchant, Fl.Au. 8:10 (1982)

nom.illeg.

At generic level, *Rorella* (Hall.ex) Allioni is a superfluous name for ***Drosera***, originally containing only *R.longifolia* (= ***D.anglica***) and *R.rotundifolia* (= ***D.rotundifolia***).

As defined by De Candolle, *sect.Rorella* was not assigned a type species, but he evidently considered this to be the typical section (including ***D.rotundifolia***).

Even when elevated to *D.subgen.Rorella* by Diels, it contained ***D.rotundifolia*** (in a separate *sect.Rossolis*, which is a superfluous name for ***sect.Drosera***). Thus, *sect.Rorella* DC., and *subgen.Rorella* (DC.) Diels are superfluous names for ***sect.Drosera*** and ***subgen.Drosera***, respectively.

Selecting a name for a new subgenus including ***D.pygmaea*** as the type species and excluding ***D.rotundifolia***, N.Merchant chose the name *Rorella*, which included ***D.rotundifolia*** in all of its circumscriptions (v.s.). Violating the original intention of Diels (*Rorella* should include ***D.rotundifolia***), this is considered illegitimate. For these reasons, it is proposed here to elevate the rank of Planchon's *sect.Bryastrum*, and thus to retain the type of N.Merchant's *subgen.Rorella*, but replace it with a legitimate name.

When elevated to subgenus (because of style morphology and the unique formation of asexual propagules called gemmae), this purely SW Australian (*sect.Lamprolepis*) or SE Australian to New Zealandic group (*sect.Bryastrum*, monotypic) group includes two subsets, separated from each other geographically as well as morphologically.

Even if the value of sectional distinction has been doubted in recent times (cf. Cheek, 1990), it is considered necessary in the context of the present grouping (especially compared with the other sections recognized here).

D.subgen.Bryastrum sect.Bryastrum Planch., l.c.:94

S: *D.sect.Rorella* DC., l.c. p.p.

D.sect.Rorella auct. non DC.: N.Merchant, l.c.

D.subgen.Bryastrum sect.Lamprolepis Planch., l.c.:93

T: ***D.platystigma*** Lehm.

S: *D.sect.Rorella* DC., l.c. p.p.

D.sect.Bryastrum auct. non Planch.: Seine & Barthlott, l.c.:585 p.p.

D.subgen.Phycopsis (Planch.) Schlauer, *stat. nov.*

BN: *D.sect.Phycopsis* Planch., Ann.sci.nat.3.ser.9:93 (1848)

T: ***D.binata*** Labill.

S: *D.sect.Ergaleium* DC., l.c. :319 p.p.

A monotypic subgenus (from E Australia to New Zealand) intermediate between the previous subgenera and those below, but closer to the last (by phytochemistry, style morphology). It is unique at first glance (forked laminal!), and it cannot be united with any of these.

D.subgen.Ergaleium (DC.) Drude, l.c. :271

BN: *D.sect.Ergaleium* DC., Prodr.1:319 (1824)

T: ***D.menziesii*** R.Br. ex DC.

The most natural grouping of all recognized here, sharing (apparently with one notable exception) corm formation and basally multipartite style branches, and almost

endemic to Australia, only two species reaching New Zealand (*D.peltata* ssp.*auriculata*) tropical Asia (*D.p.ssp.peltata*) and even E Africa (*D.insolita*, almost certainly a recent-synanthropous?-range extension). The systematic tripartition by DeBuhr (1977) has not been doubted since.

D.subgen.Ergaleium sect.Ergaleium

- S: *D.sect.Polypeltes* Diels, (1906):62 p.p. nom.superfl.
T: ***D.menziesii*** R.Br. ex DC.
D.ser.Scutelliferae Planch., l.c.:95 nom.superfl.
T: ***D.menziesii*** R.Br. ex DC.
D.ser.Luniferae Planch., l.c.
T: ***D.peltata*** Thunb.

D.subgen.Ergaleium sect.Stolonifera (Planch.) DeBuhr, Austral.J.Bot.25:215(1977)

- BN: *D.subser.stoloniferae* Planch., Ann.sci.nat.3.ser.9:95 (1848)
T: ***D.stolonifera*** Endl.

D.subgen.Ergaleium sect.Erythrorhiza (Planch.) Diels, (1906):62

- BN: *D.ser.erythrorhizae* Planch., Ann.sci.nat.3.ser.9:95 (1848)
T: ***D.erythrorhiza*** Lindley
S: *D.ser.rosulatae* Lehm., Pugill.8:36 (1844)
T: ***D.rosulata*** Lehm.
D.subser.rosulatae (Lehm.) Planch., l.c.
BN: *D.ser.rosulatae* Lehm.

New combinations in ***Drosera*** L.

In dealing with the whole genus on a world wide basis, the circumscriptions of ranks should preferably be comparable to each other. Theoretical as well as practical reasons necessitate alterations of rank in several taxa. For infraspecific subdivision as proposed here, subspecies are allopatric, whereas varieties are sympatric. The rank of form does not seem applicable in a genus as variable as ***Drosera***.

D.barbigera* subsp.*silvicola (Lowrie & Carlquist) Schlauer comb. & stat. nov.

- BN: *D.silvicola* Lowrie & Carlquist, Phytologia 73:105 (1992)
T: 7 km S N Bannister on the Albany Hwy., W.A., 11. 11. 1991, A.Lowrie 513 (PERTH)

D.citrina* var.*nivea (Lowrie & Carlquist) Schlauer comb. & stat.nov.

- BN: *D.nivea* Lowrie & Carlquist, Phytologia 73:104 (1992)
T: beside Midlands Rd., 37.3 km SE Carnamah, ca. 10 km SE of Coorow, W. A., 22. 9. 1990, A.Lowrie 278 (PERTH)

D.dichrosepala* subsp.*enodes (N.Merchant & Lowrie) Schlauer comb. & stat.nov.

- BN: *D.enodes* N.Merchant & Lowrie, Kew Bull.47:323 (1992)
T: NE Augusta, junction of Courtney Road and Scott River Road, W.A., 9. 11. 1983, A.Lowrie 83/049 (PERTH)

D.parvula* subsp.*sargentii (Lowrie & N.Merchant) Schlauer comb. & stat. nov.

- BN: *D.sargentii* Lowrie & N.Merchant, Nuytsia 8:330 (1992)
T: Junction of Stockyard Road & Merivale Road, SE corner, c. 20 km E Esperance, W.A., 22. 11. 1989, A.Lowrie s.n. (PERTH)

D.paleacea* subsp.*stelliflora (Lowrie & Carlquist) Schlauer comb. & stat. nov.

- BN: *D.stelliflora* Lowrie & Carlquist, Phytologia 73:107 (1992)

T: at motocross track, E end of N Jindong Rd., S Busselton, W.A., 24. 11. 1990,
A.Lowrie 204 (PERTH)

D.paleacea subsp.leioblastus (N.Merchant & Lowrie) Schlauer comb. & stat.nov.

BN: *D.leioblastus* N.Merchant & Lowrie, Kew Bull.47:325 (1992)

T: Brand Highway, 14.3 km NW Cataby, W.A., 29. 9. 1985, A.Lowrie 85/084
(PERTH)

D.paleacea subsp.roseana (N.Merchant & Lowrie) Schlauer comb. & stat.nov.

BN: *D.roseana* N.Merchant & Lowrie, Kew Bull.47:327 (1992)

T: Millbrook Road, 5 km E Albany Highway, W.A., 7. 10. 1987, A.Lowrie 87/025
(PERTH)

D.occidentalis var.microscapa (Debbert) Schlauer comb. & stat.nov.

BN: *D.microscapa* Debbert, Mitt.Bot.Staatss.Muenchen 30:377 (1991)

T: S coast of W.A., P.Debbert 94 (M)

D.nitidula var.allantostigma (N.Merchant & Lowrie) Schlauer stat. nov.

BN: *D.nitidula subsp.allantostigma* N.Merchant & Lowrie, Kew Bull.47:325
(1992)

T: Brand Highway, 1.3 km N Hill River, W.A., 7. 11. 1987, A.Lowrie 87/056
(PERTH)

D.nitidula var.leucostigma N.Merchant & Lowrie) Schlauer stat. nov.

BN: *D.nitidula subsp.leucostigma* N.Merchant & Lowrie, Kew Bull.47:325 (1992)

T: Brand Highway, 14.3 km NW Cataby, W.A., 7. 11. 1987, A.Lowrie 87/058
(PERTH)

D.gigantea var.geniculata (N.Merchant & Lowrie) Schlauer stat. nov.

BN: *D.gigantea subsp.geniculata* N.Merchant & Lowrie, Kew Bull.47:316 (1992)

T: 2 km N Brennans Ford on Scott River Road, W.A., 16. 9. 1984, A.Lowrie s.n.
(PERTH)

D.stricticaulis subsp.eremaea (N.Merchant & Lowrie) Schlauer comb.nov.

BN: *D.macrantha subsp.eremaea* N.Merchant & Lowrie, Kew Bull.47:318 (1992)

T: 30 km S Mt. Magnet, W.A., 1. 7. 1984, A.Lowrie 84/072 (PERTH)

One apparently new variety of **D.nitidula** (confused with **D.n.subsp.omissa** by A.Lowrie in Carniv. Pl. of Australia 2,1989) cannot be given a valid name at present because no specimen was cited. This is symbolized by "var.?" in the key.

The "collective species" **D.capillaris**, **D.montana** and **D.leucoblastia** need a reinvestigation. As long as this is not performed, it is assumed best to maintain the names and ranks which were used as the "microspecies" were first described (with some exceptions in **D.montana**). Also, a thorough examination of the widespread and polymorphic species **D.spatulata** may allow infraspecific subdivision in the future.

A perhaps unusual type of numbering is presented with this key. The position of a digit in a number reflects the position of a corresponding character pair in the key. The value of a digit reflects the character state. Thus, the opposite of

"001100. Lamina lanceolate, 4-7 mm long"

is

"001101. Lamina ovoid or circular, up to 3 mm long"

and vice versa. This kind of numbering has the advantage of facilitating direct comparison and identification of the difference between any two given taxa by just comparing their appropriate numbers, e.g.:

"1011011. (...)
D.macrantha Endl."

and

(**D.stricticaulis**...)

"1010101. (...) **subsp. eremaea** (N.Marchant & Lowrie) comb.nov."

The most significant different digit (in this case at 4. position) indicates the specific difference between the two, i.e.:

"1011. Styles divided to base but not apically"

vs.

"1010. Styles divided to base and apically plurifid"

Key to the genus **Drosera** L.

0. Plants without corms, stipules present or styles basally bipartite or entire, leaves never peltate

00. Lamina entire, never dichotomously branched, styles entire or dichotomously branched and not basally multipartite

000. Styles connate or divided or at least stigmatic apex flabellately multifid or dichotomously branched, no asexual propagules (gemmae) formed

0000. Gynoecium 5-merous

Subgen. Thelocalyx (Planchon) stat.nov.

00000. Leaves obovate, longer than 12 mm, stigmata dichotomously divided

D.sessilifolia St.Hil.

00001. Leaves cuneate, shorter than 12 mm, stigmata flabellately multifid

D.burmannii Vahl

0001. Gynoecium 3-merous

00010. Leaves exstipulate, leafbase sheathing, flowers single, rarely 2-3, pedicels glabrous, petals not upturning and uniting after anthesis

Subgen. Arcturia (Planch.) stat. nov.

000100. Leaves linear, lamina continuous with petiole, sepals oblong, not much shorter than petals

D.arcturi Hook.

000101. Lamina spathulate, sepals scarcely longer than wide, much shorter than petals

D.stenopetala Hook.f.

00011. Leaves not sheathing, petals upturning and uniting after anthesis

000110. Styles connate for at least 1/2 of their length
- Subgen. Stelogyne* (Diels) stat. nov.
- D. hamiltonii* C.R.P.Andrews
000111. Styles not connate, divergent from base
000110. Styles not divided below stigmatic area
0001100. Flowers single, stem branched, stipules present, leaves petiolate, lamina oblanceolate and up to 5 cm long
- Subgen. Meristocaulis* (Maguire & Wurdack) stat. nov.
- D. meristocaulis* Maguire & Wurdack
0001101. Flowers mostly not single, stem not branched, stipules absent, petioles indistinct, lamina linear, usu. longer than 20 cm
- Subgen. Regiae* Seine & Barthlott
- D. regia* Stephens
0001111. Styles divided below stigmatic area
0001110. Floral bracts absent, inflorescence many-flowered, styles repeatedly dichotomously divided
000111100. Scapes glandular, lamina not subpeltate-peltate
- Subgen. Coelophylla* (Planch.) stat. nov.
- D. glanduligera* Lehm.
000111101. Scapes eglandular pubescent, lamina subpeltate-peltate
- Subgen. Lasiocephala* (Planch.) stat. nov.
000111102. Stems elongate, ascending
- D. banksii* R.Br.
000111103. Stems short, acaulescent
000111104. Tomentum consisting of single or shortly branched hairs
000111105. Petioles narrower than 5 mm
000111106. Stigma branched from centre, petioles broader than 1 mm in the broadest point
- D. dilatatopetiolaris* Kondo
000111107. Stigma branched from base, petioles up to 1 mm wide
- D. petiolaris* R.Br.ex DC.
000111108. Petioles broader than 5 mm
- D. falconeri* Tsang ex Kondo
000111109. Tomentum of dendritic hairs, woolly
000111110. Petiole linear with a maximum width 1-1.5 mm; lamina 2-2.5 mm long, 2.5-3 mm wide; pedicels 1.5-2.5 mm long
- D. lanata* Kondo
000111111. Petiole oblanceolate with a maximum width 2-4 mm; lamina 3-4 mm long, 3.5-5 mm wide; pedicels 2-4.5 mm long
- D. ordensis* Lowrie
000111112. Floral bracts present or inflorescence single flowered
- Subgen. Drosera*
000111113. Seeds ovoid or ellipsoid, testa not produced beyond embryo
000111114. Stipules absent, reduced to two lateral trichomes or rarely present, if stipules present leaves with short indistinct petioles and large, broad laminae and anther thecae separated by large connective and petals scarcely reaching sepals in length
000111115. Anther thecae separated by large connective, petals scarcely reaching sepals in length
- Sect. Prolifera* C.White
000111116. Leaves with very short, indistinct petioles, lamina longer than wide, stipules conspicuous
000111117. Lamina pointed apically, broadest near the centre
- D. adelae* F.Muell.
000111118. Lamina truncate to emarginate apically, broadest near apex

D.schizandra Diels

000111110001. Leaves distinctly petiolate, lamina reniform, stipules reduced to lateral, gland-tipped trichomes

D.prolifera C.White

00011111001. Anther thecae not separated by large connective, petals longer than sepals

000111110010. Inflorescence arising laterally from ascending stem and bearing usu. more than 6 flowers, roots not thickened as storage organs

Sect.Arachnopus Planch.

D.indica L.

000111110011. Inflorescence arising centrally from basal rosette or terminal on ascending stem with up to 4 flowers, roots thickened as storage organs if stems ascending

Sect.Ptycnostigma Diels

0001111100110. Stipules present, adnate for their whole length with the exception of two inconspicuous lateral setae, caulin leaves absent

00011111001100. Petals shorter than 8 mm or scape absent

000111110011000. Scape conspicuous

0001111100110000. Seeds papillate

D.brevifolia Pursh

0001111100110001. Seeds favose

00011111001100010. Leaves distinctly petiolate, lamina spathulate-orbicular, scape glabrous

D.uniflora Willd.

00011111001100011. Leaves sessile, lamina cuneate, scape glandular-pilose

D.trinervia Spreng.

000111110011001. Scape absent, flowers almost sessile

D.acaulis L.f.

00011111001101. Petals longer than 10 mm, scape conspicuous

D.pauciflora Banks ex DC.

0001111100111. Stipules absent, caulin leaves present

D.cistiflora L. s.l.

00011111001110. Petals up to 20 mm long, with a dark base

D.cistiflora L. s.str.

00011111001111. Petals up to 10 mm long, without dark base

D.alba Phill.

0001111101. Stipules always present and conspicuous, if adnate then lateral stipule-segments large and divided and leaves with long and narrow laminae and petals longer than sepals

Sect.Oosperma sect.nov.

00011111010. Seeds papillate

000111110100. Stipules adnate to petiole for their whole length, laminae more than 4 times as long as wide

0001111101010. Petioles distinct 2-5 cm long

D.linearis Goldie

00011111010101. Petioles indistinct, almost absent

D.filiformis Raf.

0001111101010110. Leaves up to 25 cm long, stamens up to 7 mm long, glands on leaves red

var.filiformis

0001111101010111. Leaves up to 50 cm long, stamens longer than 8 mm, glands on leaves green

var.tracyi (Macf.) Diels

000111110101. Stipules adnate to petiole for up to 1/2 of their length, laminae not more than 3 times as long as wide

0001111101010. Seeds crateriform-papillate, scapes erect, petiole distinct

00011111010100. Scapes glabrous or inconspicuously glandular-puberulent

000111110101000. Plants forming stems covered by remains of dead leaves

000111110101001.	D. hirticalyx R.Duno & Culham Plants with +/- flat rosette, not forming stems
0001111101010010.	D. capillaris Poir. s.lat. Scapes longer than 5 cm, sepals longer than 2 mm
0001111101010011.	D. capillaris Poir. s.str. Scapes shorter than 5 cm, sepals shorter than 2 mm
00011111010101.	D. tenella Willd. ex Roem. & Schult. Scapes with conspicuous indumentum
000111110101010.	Scapes glandular-pilose
0001111101010111.	D. panamensis Correa & A.S.Taylor Scapes eglandular pilose
00011111010111.	D. colombiana Fernandez-Perez Seeds bullate to muricate-papillate, scapes ascending, plants acaulescent and petiole gradually widening into lamina, or +/- ascending with distinct petioles
00011111010110.	Seed papillae shallow and inconspicuous (seeds almost foveolate- reticulate), scapes pubescent, mostly also glandular, petioles short, gradu- ally broadening into lamina, plants acaulescent, scapes 1-15 cm long with usu. fewer than 10 flowers,
000111110101100.	D. montana St.Hil. s.l. Scapes only inconspicuously ascending with narrow curve at the base
0001111101011000.	D. montana St.Hil. s.str. Scape base glandular-pilose
0001111101011001.	var.montana Scape base eglandular-pilose
00011111010110010.	Sepals glandular-pilose, scape apex only with +/- stalked glands or glabrous
00011111010110011.	var.tomentosa (St.Hil.) Diels Sepals eglandular-pilose, scape apex eglandular-pilose
000111110101101.	var.schwackei Diels Scapes more conspicuously ascending with wide curve at the base, scapes long eglandular-pilose at base and glandular + eglandular-pilose above
0001111101011010.	D. hirtella St.Hil. Scapes red, strikingly ascending, with shorter red hairs in lower 2/3 or all of the scape, calyx lobes frequently with some eglandular hairs, petioles glabrous or almost so, leaves less numerous, usu. red
0001111101011011.	var.hirtella Scapes yellowish green, more erect, with longer, yellowish hairs in lower 1/2 of the scape, missing or very sparse in upper portion which is mainly glandular hairy, calyx lobes with only glandular hairs, petioles hairy, leaves more numerous, deeper purple red
00011111010111.	var.lutescens St.Hil Seed papillae conspicuous, plants (mostly) ascending or forming stems, petiole distinct, scapes glandular puberulent or glabrous
000111110101110.	Plants forming stems covered with remains of dead leaves, scapes glandular puberulent, 10-25 cm long, more than 4 times as long as leaves, with usu. more than 10 flowers
000111110101111.	D. roraimae (Klotzsch ex Diels) Maguire & Laundon Plants often caulescent but not covered with remains of dead leaves, scapes glabrous, shorter and not more than twice as long as leaves, with usu. less than 10 flowers
00011111011.	D. intermedia Hayne Seeds foveolate or reticulate
000111110110.	Plants ascending, scape erect
0001111101100.	Styles apically entire

0001111101101.	<i>D. bequaertii</i> Taton Styles divided apically
000111110111.	<i>D. neocaledonica</i> Hamet Plants acaulescent, leaves in basal rosettes
0001111101110.	Scapes erect
00011111011100.	Scapes glabrous or inconspicuously glandular puberulent
000111110111000.	Sepals shorter than 3 mm
000111110111001.	<i>D. esmeraldae</i> (Steyerm.) Maguire & Wurdack Sepals longer than 3 mm
00011111011101.	<i>D. pusilla</i> H.B.K. Scapes with conspicuous indumentum
000111110111010.	Scapes glandular-pilose, petiole gradually widened into
lamina	Petioles setaceous-pilose beneath, lamina narrowly oblanceolate
0001111101110100.	
00011111011101000.	<i>D. arenicola</i> Steyerm. Scapes up to 2 cm long with up to 4 flowers
	<i>var. arenicola</i>
00011111011101001.	Scapes longer than 2 cm with more than 4 flowers
	<i>var. occidentalis</i> Maguire & Wurdack
0001111101110101.	Leaves glabrous or sparingly, i.e. not setaceous pilose beneath, lamina obovate
0001111101110111.	<i>D. cayennensis</i> Sagot ex Diels Scapes eglandular-pilose, petioles distinct, lamina rotundate- obovate to suborbicular
0001111101110110.	Flowers 2-5, peduncle longer than 2 cm
0001111101110111.	<i>D. kaieteurensis</i> Brumm.-Ding. Flowers single, rarely 2, peduncle scarcely 2.5 mm long
0001111101111.	<i>D. felix</i> Steyerm.& L.B.Smith Scapes ascending
00011111011110.	Styles thickened basally, tapering towards apex
00011111011111.	<i>D. spatulata</i> Labill. Styles thickening towards apex
000111110111110.	Scapes glabrous, stigma divided
000111110111111.	<i>D. ob lanceolata</i> Y.Z.Ruan Scapes with tomentum, stigma entire
0001111101111110.	Petiole gradually widening into lamina, flowers red
0001111101111111.	<i>D. dielsiana</i> Exell & Laundon Petiole abruptly widening into lamina, flowers white to pale pink
0001111101111110.	Scapes glandular
0001111101111111.	<i>D. burkeana</i> Planch. Scapes eglandular-pilose
000111111.	<i>D. pilosa</i> Exell & Laundon Seeds narrowly fusiform with testa produced below and above the embryo
	Sect. Drosera
000111110.	Plants acaulescent with basal rosette, if scape glandular
then petiole indistinct	
0001111100.	Scapes glabrous or eglandular pubescent, lamina orbicular to linear, not cuneate, petiole glabrous or sparsely pilose beneath
to linear, not cuneate, petiole glabrous or sparsely pilose beneath	
00011111000.	Lamina linear, longer than 1.5 cm
	<i>D. anglica</i> Huds.
000111111001.	Lamina oblong to orbicular, up to 1 cm long
0001111110010.	Lamina oblong, stipules divided to base
	<i>D. communis</i> St.Hil.
0001111110011.	Lamina orbicular, stipules divided above centre
	<i>D. rotundifolia</i> L.
00011111101.	Scapes glandular-pilose, petiole indistinct or strigose-

pilose beneath 000111111010. 0001111110100. cuneate	Styles not repeatedly forked, stigma swollen Petiole indistinct, not strigose-pilose beneath, lamina
0001111110101. 00011111101010.	<i>D.cuneifolia</i> L.f. Petiole distinct Lamina orbicular-spathulate
00011111101011.	<i>D.slackii</i> M.R.Cheek Lamina linear, about 8 times as long as wide
000111111011. 0001111110110.	<i>D.cendeensis</i> Tamayo & Croizat Styles repeatedly forked, stigma not much swollen Leaves obovate, rounded at apex, thin, flowers light red to white, rosette with 1 layer of green leaves
0001111110111. red, rosette with several layers of functional leaves	<i>D.natalensis</i> Diels Leaves cuneate, +/- triangular, coriaceous, flowers darker
00011111111. then petiole distinct	Plants caulescent, stems at least 2 cm long, if acaulescent
00011111110.	Lamina scarcely 3 times as long as wide
000111111100.	Stems very long (usu. 60-90 cm)
000111111101. 0001111111010.	<i>D.elongata</i> Exell & Laundon Stems shorter (up to 30 cm)
0001111111011. 00011111110110. leaves reflexed in age	Leaves evenly spaced on the stem
000111111101100.	<i>D.katangensis</i> Taton Leaves crowded apically on the stem
Scapes +/- erect, stipules deeply dissected	Petioles +/- densely pilose, especially on the lower surface,
000111111101101. apically	<i>D.glabripes</i> (Harv.) Stein Scapes conspicuously curved basally, stipules lacerated
00011111110111. age	<i>D.madagascariensis</i> DC. Petioles sparsely pubescent on both surfaces, leaves erect in
0001111111111. 0001111111110. tum of the stem	<i>D.affinis</i> Welw.ex Oliv. Lamina at least 4 times as long as wide Stipules divided to base, inconspicuous in the dense tomentum
0001111111111. 00011111111110. 000111111111100.	<i>D.hilaris</i> Cham.& Schlechtd. Stipules entire at least to centre Leaf lamina glabrous or pilose, not woolly beneath Scapes glabrescent
000111111111101. 0001111111111010. apically, petioles with rust-brown hairs	<i>D.humbertii</i> Exell & Laundon Scapes glandular pilose Stipules divided into subulate setaceous appendages
0001111111111011.	<i>D.ramentacea</i> Burch. ex DC. Stipules slightly cleft apically, nearly entire, petioles glabrescent, not with rust-brown indumentum
000111111111111. 00011111111110.	<i>D.capensis</i> L. Lamina with wooly indumentum beneath Lamina oblong spathulate, up to 2 cm long
0001111111111111. 0001111111111110.	<i>D.chrysolepis</i> Taub. Lamina linear, at least 3 cm long Lamina scarcely 8 cm long, scapes glabrous
00011111111111111.	<i>D.villosa</i> St.Hil. Lamina longer than 10 cm, scapes villous

D.graminifolia St.Hil.

001. Styles never divided, stigmatic apex sometimes broadened or flabellate, asexual propagules (gemmae) usually formed

Subgen.Bryastrum (Planch.) *stat. nov.*

0010. Flowers 4-merous

Sect.Bryastrum

D.pygmaea DC.

0011. Flowers 5-merous

Sect.Lamprolepis Planch.

00110. shallow Stigma confluent with style, not widest near apex, lamina

001100. Lamina lanceolate, 4-7 mm long

0011000. Calyx red-hirsute, corolla orange, leaf lamina 6-7 mm long

D.barbigena Planch.

00110000. Scapes covered with long, lanate, glandular hairs, corolla red or bright orange, black in throat, styles and stigmas black

subsp.barbigena

00110001. Scapes with short, studlike glandular hairs, corolla pink, red in throat, styles red, stigmas white

subsp.silvicola (Lowrie & Carlquist) *comb. &*

stat. nov

0011001. 2.5-6 mm long Calyx transparent-hirsute, corolla white or pink, leaf lamina

D.scorpioides Planch.

001101. Lamina ovoid or circular, up to 3 mm long

0011010. Petals emarginate, 3 mm long

Deneabba N.Merchant & Lowrie

0011011. Petals ovate or longer than 3 mm

00110110. Styles 5, filiform and horizontal, stipule-cluster angled and acute or petioles flattened and 2-2.5 mm wide

001101100. Stipule-cluster angled, acute, petioles narrower than 1 mm

D.androsacea Diels

001101101. Stipule-cluster ovoid, petioles flattened, 2-2.5 mm wide

D.pulchella Lehm.

00110111. Styles 3-4, very rarely 5, stipule-cluster not angled, petioles

up to 1 mm wide 001101110. Stipule-cluster compact and hemispherical

D.pycnoblasta Diels

001101111. Stipule-clusters tipped, not compact and hemispherical

0011011110. Petals white, mostly yellow in outer half, petiole minutely glandular on upper surface only, style+stigma filiform, not thickened

D.citrina Lowrie & Carlquist

00110111100. Flowers with petals yellow in outer half

var.citrina

00110111101. Flowers white

var.nivea (Lowrie & Carlquist) *comb. & stat.nov.*

0011011111. Flowers white, pink or orange, never yellow, petiole glandular beneath, marginally, on both surfaces, or glabrous, style+stigma slightly thickened

00110111110. Scapes 4.5-12 cm long, sepals not reflexed in flower, petals (5-) 7-10 mm long, orange, white, or pink

D.leucoblasta Benth. *s.l.*

001101111100. Fruiting pedicels all erect

0011011111000. Style+stigma tapering from base

00110111110000. Flowers orange

D.echinoblastus N.Merchant & Lowrie

00110111110001. Flowers white or pink

D.helodes N.Merchant & Lowrie

0011011111001. Style+stigma widest near centre

00110111110010.	Stipule-cluster 4 mm long, stipules multipartite
001101111100100.	Flowers orange <i>D.callistos</i> N.Marchant & Lowrie
001101111100101.	Flowers white or pink <i>D.closterostigma</i> N.Marchant & Lowrie
00110111110011.	Stipule-cluster 7 mm long, compact and smooth <i>D.leucoblasta</i> Benth. s.str.
001101111101.	Fruiting pedicels reflexed
0011011111010.	Petals pandurate
00110111110100.	Flowers orange <i>D.miniata</i> Diels
00110111110101.	Flowers white or pink <i>D.walyunga</i> N.Marchant & Lowrie
0011011111011.	Petals obovate <i>D.spilos</i> N.Marchant & Lowrie
001101111111.	Scapes up to 5 cm long or sepals reflexed in flower, petals
never orange	Petals up to 6.5 mm long, dark pink <i>D.lasiantha</i> Lowrie & Carlquist
001101111110.	Petals shorter than 6 mm, white, sometimes with pink spots Flowers usu. fewer than 15, well-spaced Sepals reflexed in flower <i>D.dichrosepala</i> Turcz.
00110111111100.	Peduncle, pedicels, and base of sepals minutely glandular
001101111111000.	<i>subsp.dichrosepala</i> Peduncle, pedicels, and sepals glabrous
puberulent	<i>subsp.enodes</i> (N.Marchant & Lowrie) comb. &
001101111111001.	<i>stat.nov.</i> Sepals not reflexed in flower
00110111111101.	Petals narrowly obovate, without pink spots
0011011111110100.	Petiole widest near center, sepals elliptic <i>D.oreopodium</i> N.Marchant & Lowrie
0011011111110101.	Petiole widest near base, sepals orbicular <i>D.grievei</i> Lowrie & N.Marchant
001101111111011.	Petals broadly obovate, with pink spots <i>D.parvula</i> Planch.
0011011111110110.	Innermost fringe of lateral lobes of stipule produced into seta up to 2 mm long, pedicels erect in fruit <i>subsp.parvula</i>
0011011111110111.	Innermost fringe of lateral lobes of stipule produced into 5 mm long seta, pedicels pendulous in fruit <i>subsp.sargentii</i> (Lowrie & N.Marchant) comb. &
<i>stat. nov.</i>	
00110111111111.	<i>subsp.paleacea</i> Dc. Flowers more than 20, crowded, seemingly in several rows
00110111111110.	Scapes minutely glandular puberulent to glabrous <i>subsp.paleacea</i>
001101111111100.	Petiole up to 5 mm long, petals obovate
001101111111101.	Petiole 10 mm long, petals lanceolate <i>subsp.stelliflora</i> (Lowrie & Carlquist) comb. &
<i>stat. nov.</i>	
001101111111111.	Scapes with hairy pubescence
0011011111111110.	Scapes covered with eglandular pubescence <i>subsp.trichocaulis</i> (Diels) N.Marchant & Lowrie
0011011111111111.	Scapes covered with glandular pubescence
0011011111111110.	Stipules 3-cleft, forming compact clusters <i>subsp.leioblastus</i> (N.Marchant & Lowrie) comb.

& stat.nov.

001101111111111.	Stipules multifid, forming loose clusters <i>subsp.roseana</i> (N.Marchant & Lowrie) comb. &
00111.	Style abruptly widened into a flattened stigma or stigma clavate and widest near apex and lamina deeply concave
001110.	Stigma clavate, not knob-like or ovate, lamina deeply concave
0011100. 10 flowers	Styles 3, lamina elliptic, inflorescence with usu. more than
0011101. 10 flowers	<i>D.rechingeri</i> Strid Styles 5, lamina circular, inflorescence usu. with fewer than
00111010. 1- 2 flowers	<i>D.occidentalis</i> Morrison Rosette loosely open with 5-8 leaves, scapes c. 1 cm long with
001110100.	<i>subsp.occidentalis</i> Lateral lobes of stipules entire <i>var.occidentalis</i>
001110101.	Lateral lobes of stipules two-cleft <i>var.microscapa</i> (Debbert) comb. &
stat.nov.	
00111011. with 1-8 flowers	Rosette compact with 20-30 leaves, scapes c. 2.5 cm long
00111.	<i>subsp.australis</i> N.Marchant & Lowrie Style abruptly widened into stigma, stigma sometimes only
knob-like or ovate, lamina shallow	
0011110.	Gemmae with warts at apex, stigma ovoid or knob-like
00111100. flattened laterally	Stigma ovoid to oblong, inflorescence bracteate, gemmae
001111000.	Style and stigma dark, flowers orange <i>D.platystigma</i> Lehm.
001111001.	Style and stigma white, flowers white or pink <i>D.mannii</i> Cheek
00111101. tended dorsiventrally	Stigma knob-like, inflorescence ebracteate, gemmae flat-
001111010.	Styles 3 <i>D.hyperostigma</i> N.Marchant & Lowrie
001111011.	Styles 5 <i>D.sewelliae</i> Diels
0011111.	Gemmae with a stalked gland at apex, stigma peltate, circular or allantoid-reniform
00111110.	<i>D.nitidula</i> Planch. Stigma allantoid-reniform <i>subsp.nitidula</i>
001111100.	Leaf lamina up to 2.5 mm long, orbicular
0011111000.	Stigma reddish
00111110000.	Stigma reniform <i>var.nitidula</i>
00111110001. Lowrie) stat. nov.	Stigma allantoid <i>var.allantostigma</i> (N.Marchant &
0011111001.	Stigma white <i>var.leucostigma</i> (N.Marchant & Lowrie)
stat. nov.	
001111101.	Leaf lamina 3-5 mm long, spatulate <i>var.?</i>
00111111.	Stigma circular

subsp. omissa (Diels) N.Marchant & Lowrie
Lamina dichotomously branched, styles basally multipar-

01.
tite

Subgen. Phycopsis (Planch.) stat. nov.
D. binata Labill.

1. Plants with corms and/or leaves peltate, styles multipar-
tite, stipules always absent

Subgen. Ergaleum DC.

10. Leaves peltate, cauline, basal rosette sometimes present,
lowermost leaf whorls not fimbriate-eglandular

Sect. Ergaleum

100. Leaves usu. not 3 together, stem glabrous, sometimes
branching, sepals glabrous, if sepals not totally glabrous plants erect

1000. Sepals at least marginally deeply fringed and/or glandular
or sepals pilose on the whole surface, if sepals glabrous lamina distinctively crescentic
and plants less tan 0.5 m tall, few branched with few prophylls

10000. Leaves not crescentic, but sometimes reniform

100000. Sepals glandular throughout

D. marchantii DeBuhr

1000000. Basal bracts few, flowers pink

subsp. marchantii

1000001. Basal bracts numerous, flowers white

subsp. prophylla N.Marchant & Lowrie

100001. Sepal margins deeply fringed and/or glandular

1000010. Flowers usu. 5-merous, sepals deeply fringed and/or glan-
dular marginally

10000100. Lamina campaniform-concave, pointing downwards

D. huegelii Endl.

10000101. Lamina reniform, pointing horizontally outwards

100001010. Bracts present, inflorescence with up to 5 flowers, styles
irregularly divided

D. bulbigena C.Morrison

100001011. Bracts absent, inflorescence with more than 5 flowers,
styles divided to base into c. 18 flattened filiform segments, c. 12 horizontal and
upturning, the others erect

D. radicans N.Marchant

1000011. Flowers with usu. 8 sepals, petals, and stamens, sometimes
more, sepals marginally non-stalked glandular

D. heterophylla Lindl.

10001. Leaves distinctively crescentic

100010. Bracts dentate apically, cauline leaves rarely developed

D. insolita Taton

100011. Bracts not dentate apically, cauline leaves developed

1000110. Lamina of basal leaves transversely elliptic, flat, inflores-
cence 5-20 flowered, erect stem straight

10001100. Petioles of lower cauline leaves appressed to stem, 1-1.5 mm
long, petioles of upper cauline leaves semierect, 4-7 mm long

D. bicolor Lowrie & N.Marchant

10001101. Petioles of cauline leaves all semierect, 12 mm long

D. peltata Thunb.

100011010. Seeds usu. narrow ellipsoid 0.3-0.5 mm long, basal un-
branched part of style 0.1-0.3 mm long, sepals 2-4 mm long, hairy or glabrous, petals
5-6 mm long

subsp. peltata

100011011. Seeds narrow linear 0.5-1 mm long, basal part of style 0.3-
0.5 mm long, sepals 3-6 mm long, glabrous, petals 5- 8 mm long

subsp. auriculata (Backh.ex Planch.) Conn

1000111. Lamina of basal leaves flabellate, folded, inflorescence 1-2

flowered, erect stem flexuous

D.salina N.Merchant & Lowrie

1001. Sepals totally glabrous, if lamina crescentic plants many-branched, up to 1 m tall with many prophylls
10010. Stem usu. branching and/or inflorescence branched, sepals not iridescent green
100100. Lamina crescentic or reniform
1001000. Lamina distinctively crescentic, more than 4 mm wide, uppermost prophylls without lamina, sepals more than 2 mm long, styles up to 0.8 mm long

D.gigantea Lindl.

10010000. Leaves and lateral branches bent towards apex of branch or stem, stem erect

var.gigantea

10010001. Leaves and lateral branches sometimes bent towards base of branch or stem, stem flexuose

var.geniculata (N.Merchant & Lowrie) *stat. nov.*

1001001. Lamina broadly reniform, less than 2 mm wide, uppermost prophylls with undeveloped lamina, sepals up to 2 mm long, styles longer than 1 mm

D.graniticola N.Merchant

100101. Lamina orbicular
1001010. Inflorescence branched, petals 4-6 mm long, corm present

D.myriantha Planch.

1001011. Inflorescence not branched, petals up to 4 mm long, corm apparently absent or inconspicuous

D.subtilis N.Merchant

10011. Stem and inflorescence usu. not branched, lamina orbicular, sepals iridescent green

D.microphylla Endl.

101. Leaves usu. 3 together, stem never branching or very rarely few-branched, sepals glandular-pilose throughout, if sepals glabrous plants climbing

1010. Styles divided to base and apically plurifid

10100. Basal leaf rosette present

D.andersoniana Fitzg.ex Ewart & White

10101. Basal leaves absent

101010. Stem erect, lamina orbicular

D.stricticaulis (Diels) O.H.Sargent

1010100. Basal adventitious stolons absent

subsp.stricticaulis

1010101. Basal adventitious stolons present

subsp.eremaea (N.Merchant & Lowrie) *comb.nov.*

101011. Stem climbing or lamina crescentic

1010110. Lamina orbicular, petals yellow

D.subhirtella Planch.

10101100. Sepals hirsute

subsp.subhirtella

10101101. Sepals glabrous

subsp.moorei (Diels) N.Merchant

1010111. Lamina crescentic, petals white, pink, or pale yellow

10101110. Stem glabrous, sepals 5-7 mm long

D.neesii Lehm.

101011100. Leaves yellow green, stem up to 1.5 mm in diameter, petals pale yellow

subsp.neesii

101011101. Leaves red, stem more tan 1.5 mm in diameter, petals pink

subsp.borealis N.Merchant

10101111. Stem glandular, sepals 3-5 mm long

D.modesta Diels

1011. Styles divided to base but not apically
 10110. Petals obovate, white, corm white
 101100. Lamina reniform
- D.erythrogyne* N.Marchant & Lowrie
101101. Lamina circular
 1011010. Stem glabrous, flexuose, sepals only marginally glandular,
 ovary red
- D.pallida* Lindl.
1011011. Stem glandular pubescent, straight, sepals glandular pi-
 lose on whole surface, ovary green
- D.macrantha* Endl.
10111. Petals cuneate, red, pink or rarely white ageing pink, corm
 red or pink
- D.menziesii* R.Br.ex DC.
101110. Leaves at base of stem not crowded
 1011100. Petals not white, sepals fimbriate
 10111000. Stolon below ground up to 10 cm long, corm red
- subsp.menziesii*
10111001. Stolon below ground up to 45 cm long, corm white blushed
 pink
- subsp.penicillaris* (Benth.) N.Marchant & Lowrie
1011101. Petals white, ageing pink, sepals distally fimbriate
- subsp.thysanosepala* (Diels) N.Marchant
101111. Leaves at base of stem crowded, forming a closed cylinder
- subsp.basifolia* N.Marchant & Lowrie
11. Leaves not peltate or lowermost whorls fimbriate-eglandular
 110. Cauline leaves present
- Sect.Stolonifera* DeBuhr
1100. Lowermost whorls of leaves fimbriate-eglandular, caulin
 leaves peltate
- D.fimbriata* DeBuhr
1101. Lowermost leaves neither fimbriate nor eglandular, caulin
 leaves with the margins of petiole and lamina confluent
11010. Cauline leaves whorled, stigmas clustered in two groups,
 one erect, the other spreading horizontally
- D.stolonifera* Endl.
110100. Lamina of the upper leaves obovate or reniform
 1101000. Leaves and stems red, yellow-red, or dark green
 11010000. Lateral branches erect or absent
 110100000. Rosette leaves transversely elliptic, upper leaves with peti-
 oles up to 5 mm long
- subsp.stolonifera*
110100001. Rosette leaves spatulate, upper leaves with petioles 10-30
 mm long
1101000010. Petals pink, secondary cormiferous stolons present
- subsp.monticola* Lowrie & N.Marchant
1101000011. Petals white, secondary stolons absent
- subsp.compacta* N.Marchant & Lowrie
11010001. Lateral branches prostrate
- subsp.prostrata* N.Marchant
1101001. Leaves and stems light green or yellow-green
 11010010. Leaves light green, lamina 5-8 mm long
- subsp.rupicola* N.Marchant
11010011. Leaves yellow-green, lamina 2-4 cm long
- subsp.humilis* (Planch.) N.Marchant
110101. Lamina of the upper leaves orbicular with a wedge shaped
 incision

- subsp. porrecta* N.Marchant & Lowrie
Cauline leaves not whorled, stigmas not grouped conspicuously into 2 groups
110110. Stems usu. 2, inflorescence arising from basal rosette
D.ramellosa Lehm.
110111. Stems single, inflorescence terminal
D.platypoda Turcz.
111. Cauline leaves absent, all leaves in flat basal rosette
- Sect.*Erythrorhiza*** (Planch.) Diels
1110. Scapes many-flowered or leaves 2.5-10 cm long
11100. Scape single, cymose, erect in flower and fruit, leaves broadly spatulate or flabellate, up to 5 cm long
111000. Lamina broadly spatulate, green, red, or green with a red midrib
- D.erythrorhiza* Lindl.
1110000. Midrib of leaves not raised
11100000. Flowering after leaves are well developed
11100000. Leaf lamina broadly obovate, almost flabellate 3 cm long and wide
- subsp. erythrorhiza*
111000001. Leaf lamina obovate, elliptic or oblong, to 6 cm long
- subsp. collina* N.Marchant & Lowrie
11100001. Flowering before leaves develop
- subsp. squamosa* (Benth.) N.Marchant & Lowrie
1110001. Midrib of leaves slightly raised
- subsp. magna* N.Marchant & Lowrie
111001. Lamina flabellate, distal margin red
- D.zonaria* Planch.
11101. Scapes 3-40 or if 1 prostrate in fruit, single-flowered or with up to 6 flowers, leaves spatulate to obovate, 2-10 cm long, often reddish
111010. Scapes erect in fruit, leaves sessile
- D.macrophylla* Lindl.
1110100. Scapes 2-4-flowered
- subsp. macrophylla*
1110101. Scapes 1-flowered
- subsp. monantha* Lowrie & Carlquist
111011. Scapes prostrate in fruit, leaves petiolate
- D.prostratoscaposa* Lowrie & Carlquist
1111. Scapes single-flowered, rarely 2-flowered, leaves 0.8-3.5 cm long
11110. Petals 5-10 mm long, leaves entire
111100. Midrib of leaves raised
1111000. Styles short, tubaeform
- D.tubaestylis* N.Marchant & Lowrie
1111001. Styles filiform
11110010. Styles divided, scape erect in fruit, flowering at the end of the rainy season
- D.browniana* Lowrie & Carlquist
11110011. Styles entire, scape secund in fruit, flowering at the beginning of the rainy season
- D.bulbosa* Hook.
11110010. Calyx punctate, lamina to 2.5 cm long
- subsp. bulbosa*
11110011. Calyx not punctate, lamina up to 5.5 cm long
- subsp. major* (Diels) N.Marchant & Lowrie
111101. Midrib of leaves depressed
1111010. Leaves obovate, almost sessile
- D.rosulata* Lehm.

1111011.	Leaves spathulate or orbicular, petiolate
11110110.	Leaves spathulate, petioles gradually broadening into lamina, crowded more than 10
11110111.	<i>D.lowriei</i> N.Marchant Leaves orbicular, petioles abruptly broadened into lamina, 4-6
11111.	<i>D.orbiculata</i> N.Marchant & Lowrie Petals 10-12 mm long, leaves crenate-dentate apically
111110.	<i>D.whittakeri</i> Planch. Plants without stolons
111111.	<i>subsp.whittakeri</i> Plants with cormiferous stolons <i>subsp.aberrans</i> Lowrie & Carlquist

Acknowledgements

The directors, curators, and staff of the following herbaria have supported this work with their kind loan of specimens and communication of information:
 Tuebingen (TUB), Paris (P), Berlin (B), Munich (M), this is appreciated with sincere thanks. Special thanks to Terry Bertozzi who supplied valuable "last minute" information on *D.ordensis* when the manuscript was nearly completed (in 1994).

Note added in proof:

Thanks to Fernando Rivadavia who has informed me about the recent publication by Duno & Culham (1995), the thorough discussion of which having led to a revision (and, as is hoped, improvement!) of the key (in *sect.Oosperma*).

References

- DeBuhr, L.E. (1977) "Sectional classification of *Drosera* subgen. Ergaleium (Droseraceae)", Austral.J.Bot.25:209-218
- DeCandolle, A. (1824) "Droseraceae", Prodromus 1:317-319
- Cheek, M.R. (1990) "A new species of pygmy *Drosera* from Western Australia and a note on the status of *sect.Bryastrum* and *sect.Lamprolepis*", Phytologia 68:85-89
- Culham, A. & Gornall, R.J. (1994) "The taxonomic significance of naphthoquinones in the Droseraceae", Biochemical Systematics and Ecology 22:507-515
- Diels, L. (1906) "Droseraceae", A.Engler, Das Pflanzenreich 26
- Duno de Stefano, R. & Culham, A. (1995) "Dos Especies Nuevas del Genero *Drosera* (Droseraceae) en Venezuela y Otros Commentarios Taxonomicos" Novon 5:241-245
- Kondo, K. & Lavarack, P.S. (1984) "A cytotoxic study of some Australian species of *Drosera* L. (Droseraceae)" Bot.J.Linn.Soc.London 88:317-334
- Marchant, N. (1982) "Droseraceae", Fl.Australia 8:9-64, 383-385
- Planchon, J.-E. (1848) "Sur la famille des Droseracees", Ann.sci.nat.3.ser.9:79-99, 185-207, 285-309, pl.5 & 6
- Rivadavia, F. (1994) pers. comm.
- Schlauer, J. (1987) "Nomenclatural Synopsis of Carnivorous Phanerogamous Plants", Carnivorous Plant Newsletter 15:59-117

Seine, R. & Barthlott, W. (1994) "Some proposals on the infrageneric classification of *Drosera* L." *Taxon* 43:583-589

Takahashi, H. & Sohma, K. (1982) "Pollen Morphology of the Droseraceae and its Related Taxa", *Sci. Rep. Tohoku Univ. 4.ser. Biol.* 38:81-156

Zenk, M.H., Fürbringer, M. & Steglich, W. (1969) "Occurrence and distribution of 7-methyljuglone and plumbagin in the Droseraceae", *Phytochemistry* 8:2199-2200

International Carnivorous Plant Conference

May 16-20, 1997

hosted by

Atlanta Botanical Garden and the International Carnivorous Plant Society
Held at The Atlanta Botanical Garden, Atlanta, Georgia, USA

Proposed Agenda

Thursday, 15 May

6.30pm

Welcome buffet & late registration

Friday, 16 May

8.00am-6.30pm
day

6 Speakers from around the world featured throughout the day followed by a poster session, plant sale (only artificially propagated plants) and ABG cp collection tour

Saturday, 17 May

8.00am-5.00pm

6 Speakers throughout the day followed by ABG cp collection tour

Sunday, 18 May

8.30am-3.45pm

6 Speakers throughout the day

Monday, 19 May -

Tuesday, 20 May

Two day field trip to visit cps in Georgia and surrounding states

Registration

Registration will cover welcome buffet, continental breakfast, breaks and box lunch and drink throughout the conference

*Registration **\$65.00**

*Field Trip (limited to 50 places) **\$120.00**(inclusive of accommodation, travel and food)

The conference organizers are not responsible for securing plant import/export permits and all delegates are responsible for travel, medical, and car rental insurance.

Please address inquiries to:

International Carnivorous Plant Conference,
c/o Steve Baker, Rt. 1, Box 540-19AB, Conover, NC 28613,
Day: 704-322-2050 9am - 5pm, Evening: 704-256-7035 7pm - 10pm